

THE FORMATION AND FRAGMENTATION OF PRIMORDIAL MOLECULAR CLOUDS

Tom Abel^{1,2}, Greg Bryan^{2,3} and Michael L. Norman^{2,4}

¹*Max-Planck-Institut für Astrophysik, D-85740 Garching*

²*LCA, NCSA, University of Illinois, US-61801 Urbana/Champaign*

³*Massachusetts Institute of Technology, MA, US-02139 Cambridge*

⁴*Astronomy Department, University of Illinois, Urbana/Champaign*

ABSTRACT. Many questions in physical cosmology regarding the thermal history of the intergalactic medium, chemical enrichment, reionization, etc. are thought to be intimately related to the nature and evolution of pregalactic structure. In particular the efficiency of primordial star formation and the primordial IMF are of special interest. We present results from high resolution three-dimensional adaptive mesh refinement simulations that follow the collapse of primordial molecular clouds and their subsequent fragmentation within a cosmologically representative volume. Comoving scales from 128 kpc down to 0.5 pc are followed accurately. Dark matter dynamics, hydrodynamics and all relevant chemical and radiative processes (cooling) are followed self-consistently for a cluster normalized CDM structure formation model. Primordial molecular clouds with $\sim 10^5$ solar masses are assembled by mergers of multiple objects that have formed hydrogen molecules in the gas phase with a fractional abundance of $\lesssim 10^{-4}$. As the subclumps merge cooling decreases the temperature to ~ 200 Kelvin in multiple “cold pockets” of the merger product. Within these cold pockets, quasi-hydrostatically contracting cores with masses $\sim 100 M_\odot$ and number densities $\gtrsim 10^5 \text{ cm}^{-3}$ are found. We find that less than 1% of the primordial gas in such small scale structures cools and collapses to sufficiently high densities to be available for primordial star formation. Furthermore, our results indicate that the formation of very massive objects, massive black holes, fragmentation of a large fraction of baryons into brown dwarfs or Jupiter size fragments seems, in contrast to various claims in the literature, very unlikely. The expected escape fraction of UV photons with $h\nu < 11 \text{ eV}$ is very small.

1 Introduction

Saslaw and Zipoy (1967) realized the importance of gas phase H_2 molecule formation in primordial gas for the formation of proto-galactic objects. Employing this mechanism in Jeans unstable clouds, Peebles and Dicke (1968) formulated their model for the formation of primordial globular clusters. Further pioneering studies in this subject were carried out by Takeda *et al.* (1969), Matsuda *et al.* (1969), and Hirasawa *et al.* (1969) who followed in detail the gas kinetics in collapsing objects and studied the possible formation of very massive objects (VMO’s). Hutchins (1976) then looked in greater detail at the effects of rotation and asked what minimum Jeans mass can be reached in a collapsing primordial gas cloud. In the 1980’s major contributions to

this field were made by Rees and Kashlinsky (1983), Carr *et al.* (1984), and Couchman and Rees (1986), who all studied the possible cosmological consequences of population III star formation. Their main conclusion was that for hierarchical structure formation scenarios the first objects might reheat and reionize the universe and thus raise the Jeans mass, influencing subsequent structure formation quite dramatically. Massive Pop III stars would also pre-enrich the intergalactic medium with metals.

It is clear that the uncertainties mentioned above arise due to the inherently multidimensional, nonlinear, nonequilibrium physics which determine the collapse and possible fragmentation of gravitationally and thermally unstable primordial gas clouds. The computational expense of solving the network of chem-

ical rate equations forced early studies to focus on single cell calculations (cf. Hirasawa 1969; Hutchins 1976; Palla *et al.* 1983; MacLow and Shull 1986; Puy *et al.* 1996; Tegmark *et al.* 1997) adopting simple collapse models. Bodenheimer (1986) was the first to address the hydrodynamic aspects of the problem, in spherical symmetry. Similarly, using a spherical Lagrangian hydrodynamics code and solving the kinetic rate equations simultaneously, Haiman, Thoul and Loeb (1996) studied the important question of which mass scales are able to cool efficiently enough to collapse. Unfortunately, the issue of fragmentation cannot reliably be addressed in such spherically symmetric models.

Multi-dimensional studies of first structure formation have only recently become computationally feasible (Abel 1995, Anninos & Norman 1996, Zhang *et al.* 1997, Gnedin & Ostriker 1997, Abel *et al.* 1998a, Abel *et al.* 1998b). Specifically in the context of CDM-type structure formation models we have studied the collapse of high- σ density fluctuations on small mass scales (Abel *et al.* 1998, hereafter AANZ98).

Here we present first results from three-dimensional adaptive mesh cosmological hydrodynamics following the collapse and fragmentation of the very first objects formed in hierarchical, CDM-like models of structure formation.

2 Simulations

The three dimensional adaptive mesh refinement calculations presented here use for the hydrodynamic portion an algorithm very similar to the one described by Berger and Collela (1989). The code utilizes an adaptive hierarchy of grid patches at various levels of resolution. Each rectangular grid patch covers some region of space in its parent grid needing higher resolution, and may itself become the parent grid to an even higher resolution child grid. The general implementation of AMR places no restriction on the number of grids at a given level of refinement, or the number of levels of refinement. Additionally the dark matter is followed with methods similar to the

ones presented by Couchman (1991). Furthermore, the algorithm of Anninos *et al.* (1997) to solve the accurate time-dependent chemistry and cooling model for primordial gas of Abel *et al.* (1997). Detailed description of the code are given in Bryan & Norman (1997,1998), and Norman & Bryan (1997,1998).

The simulations are initialized at redshift 100 with density perturbations of a Λ CDM model with $\Omega_B = 0.06$, $h = 0.5$, and $\sigma_8 = 0.7$. The abundances of the 9 chemical species (H, H^+ , H^- , He, He^+ , He^{++} , H_2 , H_2^+ , e^-) and the temperature are initialized as discussed in Anninos and Norman (1996). After a collapsing high- σ peaks has been identified in a low resolution simulation the simulations is reinitialized with multiple refinement levels on the Lagrangian volume of the collapsing structure. The mass resolution in the initial conditions within this Lagrangian volume are $0.53 M_\odot$ in the gas and $8.96 M_\odot$ for the dark matter component. The refinement criterion ensures the local Jeans length to be resolved by at least 4 grid zones as well as that no cell contains more than 4 times its initial mass of $0.53 M_\odot$. We limit the refinement to 12 levels within a 64^3 top grid which translates to a maximum dynamical range of $64 \times 2^{12} = 262,144$. As we will show below the simulation is not resolution but physics limited.

3 Results

In Figure 1 we show mass weighted, spherically averaged quantities around the densest cell found in the simulation at redshift 19.1 of various interesting quantities. In panel c) we indicate two distinct regions. Region I) ranges from the virial radius to $r_{cool} \sim 5$ pc, the radius at which the infalling material has cooled down to $T \sim 200$ K. For most of this region the H_2 cooling time t_{H_2} is shorter than the free-fall time, $t_{ff} = [3\pi/(32G\rho)]^{1/2}$, as is illustrated in panel b) of Figure 1. The H_2 number fraction rises from 7×10^{-6} to 2×10^{-4} as the free electron fraction drops from 2×10^{-4} to 2×10^{-5} (panel a).

At r_{cool} the free-fall time becomes smaller than the cooling time. Also




Figure 1. *Spherically averaged mass weighted profiles around the baryon density peak at $z=19.1$. Panel a) shows the molecular hydrogen number fraction f_{H_2} and the free electron number fraction x . Panel b) plots the H_2 cooling time t_{H_2} , the time it takes a sound wave to travel to the center, t_{cross} , and the free-fall time $t_{ff} = [3\pi/(32G\rho)]^{1/2}$. Panel c) and d) show the temperature, baryonic number density and enclosed gas mass as a function of radius. The vertical dotted line indicates the cooling radius ($\sim 5pc$). The virial radius of the $5.6 \times 10^6 M_\odot$ halo is 106 pc. The cell size on the finest grid corresponds to 0.024 pc. Note that the simulation box size corresponds to 6.4 proper kpc.*

at r_{cool} the time it takes a sound wave to travel to the center, $t_{cross} = r/c_s = 7.6 \times 10^6 r_{pc}/\sqrt{T_K} \text{ yrs}$, becomes shorter than the cooling time. The cooling time approaches a constant value at small radii (high densities) due to the transition from non-LTE to LTE populations of the H_2 rotational/vibrational states.

From the time scales one would conclude that below r_{cool} the gas should evolve quasi-hydrostatically on the cooling time scale. This is, however, not strictly true. It turns out that centrifugal forces play an

important role in the collapse. To illustrate this Figure 2 shows the mass weighted radial profiles of various energies and force terms. From the lower panel of Figure 2 it is evident that pressure + centrifugal forces dominate gravity in the range of $0.3 < r/1 \text{ pc} < 10$ yielding a net outward force (deceleration). Indeed the gas has zero radial velocity at $r = 0.3 \text{ pc}$, separating the contracting core of $\sim 100 M_\odot$ at smaller radii. There are multiple possible origins for a “centrifugal barrier” as the one observed at $r = 0.3 \text{ pc}$. The simplest explanation being angular momentum conservation. However, this would imply $v_\perp \times r = \text{const.}$ and the centrifugal force, F_c , would be proportional to r^{-3} . However, $F_c \propto r^{-1}$ holds for more than three orders of magnitude, as can be seen from Figure 2. However, the cooling instability and the continuous merging of small scale structure offer simple alternative explanations.

4 Discussion

Multiple interesting features of the collapsing and fragmenting “primordial molecular cloud” are identified. Most notably is the fact of the existence of a contracting quasi-hydrostatic core of $\sim 100 M_\odot$. Within this core the number densities increase from 10^5 to 10^8 cm^{-3} . For densities $\gtrsim 10^6 \text{ cm}^{-3}$, however, three-body formation of H_2 becomes the dominant formation mechanism transforming all hydrogen into its molecular form (Palla *et al.* 1983). Our chemical reaction network does not include this reaction and the solution cannot be correct at $r \lesssim 0.1 \text{ pc}$. The most interesting effect of the three-body reaction is that it will increase the cooling rate by a factor $\sim 10^3$ leading to a further dramatic density enhancement within the core. This will decrease the dynamical timescales to $\ll 100$ years effectively “decoupling” the evolution of the fragment from the evolution of its “host primordial molecular cloud”. Silk (1983) has argued that (due the enhanced cooling from the 3-body produced H_2) fragmentation of this core might continue until individual fragments are opacity limited (i.e. they become




Figure 2. Spherically averaged mass weighted profiles around the baryon density peak at $z=19.1$. The upper panel compares local gravitational, kinetic, and thermal energies. The lower panel compares centrifugal ($[(v)_{rms}^2 - \langle v_r \rangle_{rms}^2]/r$), pressure ($\rho^{-1} \partial P / \partial r$), and gravity ($GM(r)/r^2$) forces. They are multiplied by rm_H/k_B for illustration purposes. Within 0.3 pc gravitation dominates leading to a steeper density profile than in the region from 0.3 to 10 pc where the sum of centrifugal and pressure forces dominate.

opaque to their cooling radiation). Verifying this possibility will have to await yet higher resolution simulations.

Clearly, it is the evolution of the $\sim 100 M_\odot$ core that could participate in population III star formation. If it would form an open star cluster with 100% efficiency about 6×10^{63} UV photons would be liberated during the average life time of massive star ($\sim 5 \times 10^7$ years). This is about hundred times more than the $\sim 4 \times 10^{61}$ hydrogen atoms within the virial radius. However, the average recombination time $(nk_{rec})^{-1} \sim 5 \times 10^5$ years within the virial radius is a factor 100 less than the average lifetime of a massive star. Hence, one expects only very small or zero UV escape fractions for these objects.

The H_2 column density from the core to the virial radius is $\sim 10^{20} \text{ cm}^{-2}$. This is much larger than the typical 10^{14} cm^{-2} required for self-shielding in stationary photodissociation regions (cf. Bertoldi and Draine 1996). Hence also many sub-Lyman limit photons will be absorbed locally and cannot reach the IGM. The host primordial molecular cloud is only transparent for pho-

tons below $\sim 11 \text{ eV}$ where no Lyman Werner Band absorption can occur.

These optical depth arguments will almost certainly become incorrect once supernova explosion alter the hydrodynamics and the chemistry and cooling via collisional ionization, dissociation and shock heating. More detailed understanding of the role of local feedback will have to await new more detailed simulations.

5 Conclusions

We have reported first results from an ongoing project that studies the physics of fragmentation and primordial star formation in the cosmological context. The discussed results clearly illustrate the advantages and power of structured adaptive mesh refinement cosmological hydrodynamic methods to cover a wide range of mass, length and timescales reliably. All findings of AANZ98 are confirmed in this study. Among other things, these are that

a significant number fraction of hydrogen molecules is only found in structures at the intersection of filaments

only a few percent of the gas in a virialized halo are cooled to $T \ll T_{\text{vir}}$.

The improvement of a factor ~ 1000 in resolution over AANZ98 has given new insights in the details of the fragmentation process and constraints on the possible nature of the first structures:

Only $\lesssim 1\%$ of the baryons within a virialized object can participate in population III star formation.

The formation of super massive black holes or very massive objects in small halos seem very unlikely.

If the gas were able to fragment further through 3-body and/or opacity limited fragmentation only a small fraction of the gas will be converted into small mass objects.

The escape fraction of UV photons above the Lyman limit or in the Lyman Werner band should be small due to the high column densities of HI ($N_{\text{HI}} \sim 10^{23} \text{ cm}^{-2}$) and H₂ ($N_{\text{H}_2} \sim 10^{20} \text{ cm}^{-2}$) in the surrounding of star forming regions.

Cautionary Remark: These latter conclusions are drawn from one simulation and are to be understood as preliminary results from an ongoing investigation.

Acknowledgments

Tom Abel acknowledges support from NASA grant NAG5-3923 and useful discussions with Karsten Jedamzik, Martin Rees, and Simon White.

References

- Abel, T. 1995, Thesis, University of Regensburg, Germany
- Abel, T., Anninos, P., Zhang, Y., Norman, M. 1997a, *NewA*, 2, 181
- Abel, T., Anninos, P., Norman, M., Zhang, Y. 1998a, *ApJ*, in press
- Abel, T., Stebbins, A., Anninos, P., Norman, M.L. 1998b, *ApJ*, in press
- Anninos, P., Norman, M.L. 1996, *ApJ*, 460, 556
- Anninos, P., Zhang, Y., Abel, T., Norman, M.L. 1997, *NewA*, 2, 209
- Berger, M.J., Collela, P. 1989, *J. Comp. Phys.*, 82, 64
- Bertoldi, F., Draine, B. T. 1996, *ApJ*, 458, 222
- Bodenheimer, P.H. 1986, *Final Technical Report, California Univ., Santa Cruz*.
- Bryan, G.L., Norman, M.L. 1997, in *Computational Astrophysics*, eds. D.A. Clarke and M. Fall, ASP Conference #123
- Bryan, G.L., Norman, M.L. 1998, *in preparation*
- Carr, B.J., Bond, J.R., Arnett, W.D. 1984, *ApJ*, 277, 445
- Couchman, H. 1991, *ApJL*, 368, L23
- Gnedin, N.Y., Ostriker, J.P. 1997, *ApJ*, 486, 581
- Haiman, Z., Thoul, A.A., Loeb, A. 1996, *ApJ*, 464, 523
- Hirasawa, T. 1969, *Progr. Theoret. Phys.*, 42, 523
- Hutchins, J.B. 1976, *ApJ*, 205, 103
- Kashlinsky, A., Rees, M.J. 1983, *MNRAS*, 205, 955
- Mac Low, M.-M. & Shull, J.M. 1986, *ApJ*, 302, 585
- Matsuda, T., Sato, H., Takeda, H. 1969, *Progr. Theoret. Phys.*, 41, 840
- Norman, M.L., Bryan, G.L. 1997, in *Workshop on Structured Adaptive Mesh Refinement Grid Methods*, ed. N. Chrisochoides
- Norman, M.L., Bryan, G.L. 1998, in *Numerical Astrophysics 1998*, eds. S. Miyama & K. Tomisaka
- Palla, F., Salpeter, E.E., Stahler, S.W. 1983, *ApJ*, 271, 632
- Peebles, P.J.E., Dicke, R.H. 1968, *ApJ*, 154, 891
- Puy, D., Signore, M. 1996, *A&A*, 305, 371
- Silk, J. 1983, *MNRAS*, 205, 705
- Takeda, H., Sato, H., Matsuda, T. 1969, *Progr. Theoret. Phys.*, 41, 840
- Tegmark, M., Silk, J., Rees, M.J., Blanchard, A., Abel, T., Palla, F. 1997, *ApJ*, 474, 1
- Zhang, Y., Norman, M.L., Anninos, P., & Abel, T. 1997, in S.S. Holt and L.G. Mundy, eds., *Star formation, near and far*, AIP Press, New York, p329